





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 6 : C07D 487/04, A61K 31/505 // (C07D 487/04, 241:00, 235:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/19079 (43) International Publication Date: 29 May 1997 (29.05.97)</p>
<p>(21) International Application Number: PCT/IB96/01291 (22) International Filing Date: 22 November 1996 (22.11.96) (30) Priority Data: MI95A002446 24 November 1995 (24.11.95) IT (71) Applicant (for all designated States except US): BIOMEDICA FOSCAMA INDUSTRIA CHIMICO-FARMACEUTICA S.P.A. [IT/IT]; Via Morolense, 87, I-03013 Ferentino (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CECCARELLI, Stefano [IT/IT]; Via Dante Alighieri, 3, I-03100 Frosinone (IT). ZANARELLA, Sergio [IT/IT]; Via degli Abeti, 45, I-00010 Mentana (IT). ALTOBELLI, Maria [IT/IT]; Via Fosse Ardeatine, 102, I-03100 Frosinone (IT). D'ALESSANDRO, Alessandra [IT/IT]; Via S. Marciano, I-03039 Sora (IT). (74) Agent: MARSI, Graziella; Con Lor S.p.A., Via Renato Fucini, 14, I-20133 Milano (IT).</p>		<p>(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i></p>
<p>(54) Title: IMIDAZO[1,2-a]QUINOXALIN-4-AMINES ACTIVE AS ADENOSINE ANTAGONISTS, PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS THEREOF</p> <div style="text-align: center;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>There are described imidazo[1,2-a]quinoxalin-4-amines derivatives of formula (I) and salt thereof active as adenosine antagonists and a process for their preparation and pharmaceutical compositions containing them as therapeutically active compounds for psychiatric and neurological disorders of the central nervous system.</p> <div style="text-align: right;">  </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

IMIDAZO[1,2-a]QUINOXALIN-4-AMINES ACTIVE AS ADENOSINE ANTAGONISTS, PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS THEREOF.

5 DESCRIPTION

The present invention relates to the imidazo[1,2-a]quinoxalin-4-amines and their salts, that are active as antagonists of the adenosine receptors; a process for their preparation as well as the pharmaceutical compositions containing them as active ingredients that are useful in the therapy for the treatment of various psychiatric and neurological disorders of the central nervous system.

It is known that theophylline (1,3-dimethylxanthine) and caffeine (1,3,7-trimethylxanthine) are capable of antagonizing the effects of adenosine through interaction with its receptors, and that it is mainly to such a property that their central nervous system stimulant effects are to be ascribed. However the presence of pharmacologically relevant effects also at the heart, kidney and smooth muscle level has determined a serious limitation to the therapeutical use of these substances as agents for effectively treating the central nervous system diseases characterized by abnormalities in the neuronal transmission processes, such as, for example, depression and senile dementia. Moreover, their low affinity to the adenosine receptors implies that the therapeutically effective dosages are too close to those causing serious side effects at the central level too.


A series of compounds having various non-xanthine structures have exhibited, at different levels, affinity to the adenosine receptors (see for example documents EP 515,107 A2 and *J. Med. Chem.* 1991, 34, 1202), but none of them has the structure of imidazo[1,2-a]quinoxalin-4-amines.

Several derivatives of imidazo[1,2-a]quinoxaline derivatives have been described in the literature: for example in U.S. Patent N. 5,182,386 imidazoquinoxalinones are disclosed that interact with the central GABA receptors, whilst in WO 94/22865 analogous compounds (especially derivatives of 4,5-dihydro-4-oxoimidazo[1,2-a]quinoxaline-2-carboxylic acid) are described as antagonists of excitatory amino acids. In no case, however, an affinity of such substances to the adenosine receptors has been pointed out.

With the present invention, it has surprisingly been discovered that a group of imidazo[1,2-a]quinoxalin-4-amines are potent antagonists of the adenosine receptors that are active in vivo on the central nervous system at much lower dosages as compared with the compounds that are presently in general use in therapy.

The compounds of the invention might therefore exhibit a lower incidence of side effects, especially at the peripheral level.

The present invention relates therefore to a compound of the formula (I):



wherein:

R₁ is hydrogen or methyl;

R₂ is hydrogen, straight- or branched- chain (C₁-C₆)alkyl;

10 R₃ is hydrogen, straight- or branched- chain (C₁-C₆)alkyl
that is possibly substituted with OH, (C₃-C₈)cycloalkyl;
or R₂ and R₃ together form



20 wherein Z is a direct bond, or O, NR, R being a
straight- or branched- chain (C₁-C₆)alkyl;

m and n, same or different, are 1, 2 or 3;

R₄ and R₅ can be the same or different and are
hydrogen or halogen chosen from Cl, F, Br;


25 and its pharmacologically acceptable salts;

In the present invention, there are preferred the
compounds of formula (I) in which R₁ is hydrogen or
methyl, R₂ is hydrogen, R₃ is hydrogen, straight- or
branched- chain (C₁-C₆)alkyl that is possibly substituted
30 with OH, (C₃-C₆)cycloalkyl, R₄ and R₅ can be the same or
different and are hydrogen, chlorine or fluorine.



Especially preferred is the compound of formula (I) in which R_1 is methyl, R_2 is hydrogen, R_3 is cyclopentyl, R_4 and R_5 are both hydrogen, that is the compound 4-cyclopentylamino-1-methylimidazo[1,2-a]quinoxaline.

5 The salts of the compounds of formula (I) comprise the acid addition salts that can be prepared *in situ* during the final isolation and the purification or by means of a separate reaction of the free base with the suitable organic or inorganic acid chosen, for example,
 10 from hydrochloric, hydrobromic, phosphoric, methaphosphoric, nitric, sulphuric, tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic and p-toluene sulfonic acids.


An object of the present invention is also a
 15 process for the preparation of the compounds of the general formula (I). Said process comprises reacting the 2,3-dichloroquinoxaline of the formula (II)



wherein R_4 and R_5 are as defined for the compound of formula (I), with amino acetaldehyde dimethyl acetal or
 25 with propargyl amine, thereby to obtain, respectively, the compounds of the formulae (III) and (IV)




that are subsequently submitted to a cyclization reaction in an acidic medium, preferably at a temperature comprised between 50 and 120°C, to obtain the imidazo[1,2-a]quinoxalin-4(5H)-ones of formula (V)



in which R₁ is hydrogen or methyl depending on whether the starting compound was compound (III) or compound (IV), respectively, R₄ and R₅ being as defined above.

The transformation of the compounds of formula (V) into the inventive compounds having the formula (I) can be carried out following two different synthetic sequences: in the first one, the chloride of the formula (VI)



in which all the substituents are as already defined, said chloride being obtained by treatment of compound of formula (V) with POCl₃ or another chlorinating agent, is reacted with the suitable amine HNR₂R₃, in which R is hydrogen, straight- or branched- chain (C₁-C₄)alkyl; R₃ is

hydrogen, straight or branched chain (C₁-C₆)alkyl, possibly substituted with hydroxy, (C₃-C₄)cycloalkyl; or R₂ and R₃ together form




wherein Z is a direct bond, or O, NR₄, R₄ being a straight- or branched- chain (C₁-C₆)alkyl; m and n, same or different, are 1, 2 or 3.

15 As an alternative, the compound of formula (V) can be directly converted into the compounds of formula (I), by reacting it with a silylating agent, such as hexamethyl disilazane and the suitable amine HNR₁R₂ in which R₁ and R₂ are as defined in formula (I), at a

20 temperature between 80 and 180°C, possibly in the presence of a catalyst, such as ammonium sulphate.

The preparation of the compounds of formula (I) in which R₁ and R₂ are both hydrogen, can also be carried out by reacting the chloride of formula (VI) with

25 hydrazine to obtain a compound of formula (VII)




in which all the substituents are as defined above, followed by hydrogenation of this compound by conventional methods, such as, for example, hydrogen and palladium-on-carbon, or Raney nickel.

5 The preferred compound of the present invention can be prepared according to the general process described above through the formation of the compound of formula (V). Said compound is then converted to the preferred compounds of formula (I) by means of either synthetic
10 sequence shown above. For example, the especially preferred compound of formula (I) of the present invention can be prepared by treating 2,3-dichloroquinoxaline (compound of formula (II) in which R₁ and R₂ are both hydrogen) with propargylamine to obtain
15 the 2-propargylamino-3-chloroquinoxaline (compound of formula (IV) in which both R₁ and R₂ are hydrogen), followed by reacting this latter compound in an acidic medium, for example with concentrated sulphuric acid, at a temperature of between 50 and 120°C, to give 1-
20 methylimidazo[1,2-a]quinoxalin-4(5H)-one (compound of formula (V) in which R₁ is methyl and R₄ and R₅ are both hydrogen). This compound can finally be converted to 4-cyclopentylamino-1-methylimidazo[1,2-a]quinoxaline through the formation of the intermediate 4-chloro-1-methylimidazo[1,2-a]quinoxaline (compound of formula (VI)
25 in which R₁ is methyl and R₄ and R₅ are both hydrogen) by chlorination with one of known chlorinating agents at a temperature of between 50 and 150°C, followed by the reaction of the latter compound with cyclopentylamine, or

alternatively, by reaction of 1-methylimidazo[1,2-a]quinoxalin-4(5H)-one with cyclopentylamine in hexamethyl disilazane or other silylating agent at a temperature of between 80 and 180°C, possibly in the presence of a catalyst, such as ammonium sulphate.

The preparation of the compounds of formula (II), when they are not commercially available, can be carried out with methods such as the one shown below. Specifically, phenylenediamine (VIII) is reacted with dialkyl oxalate to give the 2,3-dihydroxyquinoxaline (IX), which is then subjected to a chlorinating reaction with one of the usual chlorinating agents, such as for example POCl₃, thereby obtaining the compounds (II)



20 As shown hereinafter in Examples 28 to 30, the compounds of formula (I) of the present invention are antagonists of the adenosine receptors, are active on the central nervous system and can therefore be advantageously used as active ingredients for the preparation of medicaments that are useful in therapy for the treatment of various psychiatric disorders of the central nervous system, such as the depressive syndromes of various etiology and symptomatology and mood disorders in general, bulimia nervosa, sleep disorders, obsessive-

25

compulsive disorders, phobias, panic attacks. Further indications are neurological diseases, such as pre-senile and senile dementia, the Alzheimer's

5 disease, the multiinfarctual dementias, the encephalopathies of toxic or traumatic origin, the Parkinson disease, the post-neurological deficits, the respiratory depression, the neonatal cerebral damage.

10 Besides being employed as drugs acting on the central nervous system, the compounds of the present invention and their salts could be used for the treatment of diseases of the renal system, such as acute renal failure, or of diseases of the cardiovascular system.

15 For all the above-mentioned therapeutical uses, the compounds of the present invention can be administered by oral, transdermal or transmucosal route, parenterally or rectally in formulations containing them as the active ingredients at a therapeutically effective dosage with conventional, non-toxic pharmaceutical excipients. The term parenteral as used herein comprises subcutaneous, 20 intravenous, intramuscular and intracerebroventricular injections. If the compounds of the present invention are in the form of a pharmaceutical composition, as in a preferred embodiment of the invention, the precise formulation employed will obviously depend on the 25 administration route chosen.

 The pharmaceutical compositions that are suitable for the oral administration can be for example tablets, aqueous or oily suspensions, dispersible powders or granules, hard or soft capsules, syrups or elixirs. The

compositions for the oral administration can contain one or more sweetening agents, colorants, flavouring and preserving agents that are suitable to make the pharmaceutical composition elegant and palatable.

5 The formulations for oral administration comprise tablets in which the active ingredient is admixed with non-toxic, pharmaceutically acceptable excipients. Said excipients can be inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate
10 or sodium phosphate; granulating or disgregating agents, such as wheat starch or alginic acid; binding agents, such as starch or gelatines; lubricant agents, such as magnesium stearate, stearic acid or talc.

 The tablets can be non-coated or coated with
15 conventional techniques known to a person skilled in the art in order to delay disintegration and absorption in the gastrointestinal tract, in order to achieve a sustained release action.

 The aqueous suspensions generally contain the
20 active ingredients admixed with the suitable excipients. The excipients can be suspending agents, such as sodium carboxymethyl cellulose, methyl cellulose, hydroxypropyl methyl cellulose, sodium alginate, polyvinylpyrrolidone; dispersants and wetting agents. They can also contain one
25 or more preservatives, such as ethyl and n-propyl p-hydroxybenzoate; one or more flavouring agent; one or more sweetening agents.

 The oily suspensions can be formulated by suspending the active ingredient in a vegetable or

mineral oil; they can contain sweetening agents and flavouring agents in order to make the preparation palatable.

5 The dispersible powders and granules that are suitable to the preparation of an aqueous suspension by adding water contain the active ingredient in admixture with the dispersing or wetting agent, a suspending agent and one or more preserving agents.

10 The pharmaceutical compositions of the present invention can also be in the form of a water/oil emulsion. The oily phase can consist of a vegetable or mineral oil. The emulsifying agents can be natural gums, such as acacia, or natural phosphatides, such as lecithins, or natural or synthetic fatty acid esters. The
15 syrops and the elixirs can be formulated with sweetening agents, for example glycerol, sorbitol or sucrose.

 The pharmaceutical compositions can be in the form of aqueous or oily, sterile injectable suspensions. The suspensions can be formulated with the known techniques
20 by using dispersing or wetting agents and suspending agents that are known in the art. The sterile injectable preparations can be sterile injectable solutions or suspensions in a non-toxic solvent or diluent that is suitable for the parenteral use.

25 The compounds of the present invention can also be administered by the rectal route in the form of suppositories. These compositions can be prepared by mixing the active ingredient with a suitable, non irritating excipient that is solid at room temperature

but liquid at the rectal temperature, thereby melting in rectum to release the drug. The polyethylene glycols and the cocoa butter are suitable compounds for this purpose.

5 The therapeutically or prophylactically effective amounts of a compound of the present invention will depend on a number of factors including, for example, the age and weight of the patient, the severity of the specific disease requiring the treatment, the administration route. However, an effective amount of the
10 compound of the present invention for the treatment of depressive syndromes and of senile dementia will generally be comprised in the range of 0.005-20 mg/kg of body weight per day, more frequently in the range of 0.05-2 mg/kg per day.

15 In order to better illustrate the present invention, the following examples are reported, that are in no way to be considered as limiting.

Example 1

(a) A mixture of 5.0 g of 2,3-dichloroquinoxaline
20 and 5.5 ml of aminoacetaldehyde dimethyl acetal in 75 ml of ethanol is refluxed for 4 h. After concentrating under vacuum, the resulting mixture is added with water and extracted with ethyl acetate. The organic extracts are then washed with saturated NaCl, dried and evaporated.
25 The residue is finally chromatographed on SiO₂ (eluent: CH₂Cl₂), thereby obtaining 5.0 g of 2-chloro-3-(2,2-dimethoxyethylamino)quinoxaline (IR (KBr): 3347, 2936, 1580, 1523, 1129 cm⁻¹). 4.5 g of this product are then treated with 20 ml of 48% HBr and the mixture is refluxed

for 4 h. After cooling down, the mixture is neutralized with aqueous NaOH and the resulting precipitate is filtered under vacuum and dried to obtain 3.2 g of imidazo[1,2-a]quinoxalin-4(5H)-one (m.p. >300°C)

5 (b) A mixture of 0.47 g of imidazo[1,2-a]quinoxalin-4(5H)-one and 0.42 ml of N,N-dimethylaniline in 5.6 ml of phosphorus oxychloride is refluxed for 2 h. After evaporating under vacuum the resulting mixture, the residue is taken up in chloroform and repeatedly washed
10 with water, then with saturated NaCl, then it is dried and evaporated thereby obtaining, after recrystallization from n-hexane / chloroform, 0.29 g of 4-chloroimidazo[1,2-a]quinoxaline (IR (KBr): 1456, 755 cm⁻¹).

 (c) A mixture of 0.25 g of 4-chloromidazo[1,2-
15 α]quinoxaline and 0.5 ml of hydrazine hydrate in 1.5 ml of ethanol is refluxed for 2 h. After cooling the mixture, the resulting precipitate is filtered under vacuum, washed and dried, thereby obtaining 0.22 g of 4-hydrazinoimidazo[1,2-a]quinoxaline (IR (KBr): 3308, 3237,
20 1570 cm⁻¹).

 (d) A mixture of 0.17 g of 4-hydrazinoimidazo[1,2-a]quinoxaline and 3.4 ml of Raney nickel in 20 ml of water is refluxed for 1.5 h. After cooling down, the mixture is filtered on Celite, followed by washing with
25 methanolic chloroform. The filtrate is concentrated under vacuum and extracted with ethyl acetate. The organic extracts are then washed with saturated NaCl, dried and evaporated, thereby obtaining 0.17 g of imidazo[1,2-a]quinoxalin-4-amine. M.p. (DSC) = 205.3°C (onset); IR

(KBr): 3301, 3143, 1653, 1525 cm^{-1} ;

$^1\text{H-NMR}$ (CDCl_3): δ 8.0 (1H, s), 7.8÷7.45 (3H, m), 7.45÷7.2 (2H, m), 5.8 (2H, sb) ; UV (EtOH): $\lambda_{\text{max}} = 229, 295, 331 \text{ nm}$.

Elementary analysis for $\text{C}_{10}\text{H}_8\text{N}_4$ (m.w. 184.20):

5 calcd. C 65.21, H 4.38, N 30.42%;

found C 65.48, H 4.71, N 30.10%.

Example 2

(a) A mixture of 10 g of 2,3-dichloroquinoxaline, 4.5 ml of propargylamine, 10.5 ml of triethylamine in 50 ml of ethanol is refluxed for 4 h. After evaporating under vacuum the resulting mixture, the residue is chromatographed on SiO_2 (eluent: CH_2Cl_2) thereby obtaining 7.0 g of 2-chloro-3-(propargylamino)quinoxaline (IR (KBr) : 3441, 3282, 1518 cm^{-1}). This product is then added with 10 ml of concentrated sulphuric acid and the resulting mixture is stirred at 90°C for 1 h. After cooling down and cautiously neutralizing with aqueous NaOH, the resulting precipitate is filtered under vacuum, washed, dried, decolorized and recrystallized from dimethylformamide, thereby obtaining 2.4 g of 1-methylimidazo[1,2-a]quinoxaline-4(5H)-one (m.p. >300°C).

(b) The chlorination of 1-methylimidazo[1,2-a]quinoxalin-4(5H)-one with POCl_3 is carried out according to a closely analogous procedure to that reported in example 1 (b), thereby obtaining 4-chloro-1-methylimidazo[1,2-a]quinoxaline (IR (KBr): 1486, 754 cm^{-1}).

(c) The reaction of 4-chloro-1-methylimidazo[1,2-a]quinoxaline with hydrazine hydrate is carried out

according to a strictly analogous procedure to that followed in example 1 (c), thereby obtaining 4-hydrazino-1-methylimidazo[1,2-a]quinoxaline (IR (KBr): 3298, 3250, 1564 cm^{-1}).

5 (d) The hydrogenation of 4-hydrazino-1-methylimidazo[1,2-a]quinoxaline with Raney nickel is carried out according to a strictly analogous procedure to that followed in example 1 (d), thereby obtaining 1-methylimidazo[1,2-a]quinoxalin-4-amine. m.p. (DSC) =
10 184.8°C (onset); IR (KBr): 3379, 3295, 1640, 1516 cm^{-1} ;
1H-NMR (CDCl_3): δ 8.05 (1H, dd), 7.9÷7.2 (3H, m), 7.2 (1H, s), 5.7 (2H, sb), 2.85 (3H, s); UV (EtOH): λ_{max} =
225, 270, 304, 329 nm. Elementary analysis for $\text{C}_{11}\text{H}_{10}\text{N}_4$ (m.w. 198.23):
15 calcd. C 66.65, H 5.08, N 28.26%;
found C 66.79, H 5.30, N 28.03%.

Example 3

A mixture of 2.8 g of 4-chloroimidazo[1,2-a]quinoxaline (example 1) and 9.8 ml of diethylamine in
20 40 ml of ethanol is refluxed for 4 h. After evaporating the solvent, the residue is taken up in chloroform and washed with water and saturated NaCl, then dried and evaporated, thereby obtaining 2.2 g of raw product that is subsequently chromatographed on SiO_2 (eluent: CH_2Cl_2):
25 after recrystallization from n-hexane there are obtained 1.3 g of 4-diethylaminoimidazo[1,2-a]quinoxaline. m.p. (DSC) = 91.7°C (onset); IR (KBr):
2976, 1518, 1425, cm^{-1} ; 1H-NMR (CDCl_3): δ 7.9 (1H, s), 7.7÷7.4 (3H, m), 7.35÷7.05 (2H, m), 4.15 (4H, q), 1.3

(6H,t); UV (EtOH): $\lambda_{\text{max}} = 231, 250, 293, 305, 332, 348 \text{ nm.}$

Elementary analysis for $\text{C}_{14}\text{H}_{14}\text{N}_4$ (m.w. 240.31):

calcd. C 69.97, H 6.71, N 23.31%;

found C 69.99, H 6.80, N 23.13%.

5 Example 4

By reaction of 4-chloroimidazo[1,2-a]quinoxaline (example 1) with isopropylamine according to a procedure that is similar to that followed in example 3 there is obtained 4-isopropylaminoimidazo[1,2-a]quinoxaline. m.p.

10 (DSC) = 102.7°C (onset); IR (KBr): 3230, 2966, 1559 cm^{-1} ;

$^1\text{H-NMR}$ (DMSO-d_6): δ 8.55 (1H,s), 8.2÷7.95 (1H,m), 7.6÷7.4 (2H,m), 7.4÷7.2 (3H,m), 4.5 (1H,m), 1.25 (6H,d); UV (EtOH): $\lambda_{\text{max}} = 227, 244, 285, 297, 318, 332 \text{ nm.}$

Elementary analysis for $\text{C}_{15}\text{H}_{14}\text{N}_4$ (m.w. 226.28):

15 calcd. C 69.00, H 6.23, N 24.76%;

found C 69.17, H 6.70, N 25.05%.

Example 5

By reaction of 4-chloro-1-methylimidazo[1,2-a]quinoxaline (example 2) with 1-ethylpropylamine, according to a procedure that is similar to that followed in example 3, there is obtained 4-(1-ethylpropylamino)-1-methylimidazo[1,2-a] quinoxaline. m.p. (DSC) = 75.9°C

(onset); IR (KBr): 3231, 2965, 1546 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.0 (1H,dd), 7.7 (1H,dd), 7.4÷7.1 (3H,m), 5.9 (1H,d), 4.2 (1H,m), 2.85 (3H,s), 1.65 (4H,m), 0.95 (6H,t); UV (EtOH): $\lambda_{\text{max}} = 221, 241, 268, 298, 313 \text{ nm.}$

Elementary analysis for $\text{C}_{15}\text{H}_{20}\text{N}_4$ (m.w. 268.36):

calcd. C 71.61, H 7.51, N 20.83%;

found C 71.12, H 7.53, N 20.79%.

Example 6

By reaction of 4-chloromidazo[1,2-a]quinoxaline (example 1) with ethanolamine according to a procedure that is similar to that followed in example 3 there is obtained 4-(2-hydroxyethylamino)imidazo[1,2-a] quinoxaline. m.p. (DSC) = 150.8°C (onset); IR (KBr): 3312, 1598, 1564 cm^{-1} ;

$^1\text{H-NMR}$ ($\text{CDCl}_3/\text{CD}_3\text{OD}$): δ 8.2 (1H,s), 8.1÷7.55 (2H,m), 7.55 (1H,s), 7.4÷7.2 (2H,m), 3.8 (4H,s); UV (EtOH): $\lambda_{\text{max}} = 227, 285, 297, 317, 330 \text{ nm}$.

Elementary analysis for $\text{C}_{12}\text{H}_{12}\text{N}_4$ (m.w. 228.25):

calcd. C 63.14, H 5.30, N 24.55%;

found C 63.16, H 5.40, N 24.99%.

Example 7

By reaction of 4-chloro-1-methylimidazo[1,2-a]quinoxaline (example 2) with ethanolamine, according to a procedure that is similar to that followed in example 3 there is obtained 4-(2-hydroxyethylamino)-1-methyl imidazo[1,2-a]quinoxaline. m.p. (DSC) = 173.4°C (onset); IR (KBr): 3415, 3217, 1558 cm^{-1} ; $^1\text{H-NMR}$

(CDCl_3): δ 8.0 (1H,dd), 7.7÷7.2 (3H,m), 7.2 (1H,s), 6.6 (1H,m), 5.3 (1H,sb), 3.85 (4H,m), 2.8 (3H,s); UV (EtOH): $\lambda_{\text{max}} = 225, 242, 271, 301, 314, 327 \text{ nm}$.

Elementary analysis for $\text{C}_{13}\text{H}_{14}\text{N}_4$ (m.w. 242.28):

calcd. C 64.45, H 5.82, N 23.12%;

found C 64.29, H 5.91, N 23.13%.

Example 8

By reaction of 4-chloromidazo[1,2-a]quinoxaline

(example 1) with cyclopentylamine according to a procedure that is similar to that followed in example 3 there is obtained 4-cyclopentylaminoimidazo[1,2-a]quinoxaline. m.p. (DSC) = 114.3°C (onset); IR (KBr):

5 3419, 2962, 1543 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 7.9 (1H,s), 7.8÷7.0 (5H,m), 6.2 (1H,d), 4.7 (1H,m), 2.3÷2.05 (4H,m), 2.0÷1.3 (4H,m); UV (EtOH): λ_{max} = 228, 244, 285, 297, 319, 333 nm.

Elementary analysis for $\text{C}_{15}\text{H}_{16}\text{N}_4$ (m.w. 252.32):

10 calcd. C 71.40, H 6.39, N 22.20%;

found C 71.13, H 7.30, N 22.29%.

Example 9

A mixture of 3.5 g of 1-methylimidazo[1,2-a]quinoxalin-4(5H)-one (example 2), 13 ml of hexamethyl
15 disilazane, 0.5 g of ammonium sulphate and 8.7 ml di-cyclopentylamine is stirred at 120°C in a Dean-Stark apparatus for 24 h. After concentrating the resulting mixture under vacuum, the residue is added with water and extracted with ethyl acetate. The organic extracts are
20 then washed with water and saturated NaCl, dried and evaporated, thereby obtaining 1.5 g of a raw product that is subsequently chromatographed on SiO_2 (eluent: $\text{CH}_2\text{Cl}_2/\text{AcOEt}$ 97:3). By recrystallizing from ethyl acetate, there is obtained 4-cyclopentylamino-1-methyl
25 imidazo [1,2-a]quinoxaline.

m.p. (DSC) = 167.3°C (onset); IR (KBr): 3294, 2948, 1544 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.0 (1H,dd), 7.7 (1H,dd), 7.4÷7.05 (3H,m), 6.0 (1H,d), 4.6 (1H,m), 2.8 (3H,s), 2.3÷2.05 (4H,m), 2.0÷1.4 (4H,m); UV (EtOH): λ_{max} = 225, 243, 272,

301, 316, 329 nm.

Elementary analysis for $C_{16}H_{16}N_4$ (m.w. 266.35):

calcd. C 72.15, H 6.81, N 21.03%;

found C 71.86, H 6.81, N 20.79%.

5 Example 10

By reaction of 4-chloroimidazo[1,2-a]quinoxaline (example 1) with cyclohexylamine, according to a procedure that is similar to that followed in example 3, there is obtained 4-cyclohexylaminoimidazo[1,2-a]quinoxaline. m.p. (DSC)= 94.0°C (onset); IR (KBr): 3227, 2924, 1560 cm^{-1} ; 1H -NMR ($CDCl_3$): δ 7.85 (1H,d, J=2Hz), 7.8÷7.4 (3H,m), 7.4÷7.1 (2H,m), 6.0 (1H,d), 4.2 (1H,m), 2.3÷1.95 (4H,m), 1.95÷0.9 (6H,m) ; UV (EtOH): λ_{max} = 225, 242, 283, 294, 316, 330 nm.

15 Elementary analysis for $C_{16}H_{16}N_4$ (m.w.266.35):

calcd. C 72.15, H 6.81, N 21.03%;

found C 72.25, H 7.04, N 21.19%.

Example 11

By reaction of 1-methylimidazo[1,2-a]quinoxalin-4(5H)-one (example 2) with cyclohexylamine, according to a procedure that is similar to that followed in example 9, there is obtained 4-cyclohexylamino-1-methyl imidazo [1,2-a]quinoxaline. m.p. (DSC)=126.5°C (onset); IR (KBr): 3345, 2938, 1546 cm^{-1} ; 1H -NMR ($CDCl_3$): δ 8.0 (1H,dd), 7.7 (1H,dd), 7.5÷7.1 (3H,m), 6.0 (1H,d), 4.2 (1H,m), 2.85 (3H,s), 2.3÷1.95 (4H,m), 1.95÷0.9 (6H,m); UV (EtOH): λ_{max} = 225, 243, 272, 301, 316, 329 nm.

Elementary analysis for $C_{17}H_{20}N_4$ (m.w. 280.37):

calcd. C 72.83, H 7.19, N 19.98%;

found C 72.21, H 7.54, N 20.14%.

Example 12

By reaction of 4-chloroimidazo[1,2-a]quinoxaline
5 (example 1) with piperidine, according to a procedure
that is similar to that followed in example 3, there is
obtained 4-(1-piperidinyl)imidazo[1,2-a]quinoxaline. m.p.
(DSC)=108.2°C(onset); IR(KBr): 3107, 2935, 1517 cm^{-1} ; ^1H -
NMR (CDCl_3): δ 7.9 (1H,d, J=2Hz), 7.75÷7.4 (3H,m), 7.4÷7.1
10 (2H,m), 4.3 (4H,t), 1.9÷1.6 (6H,m); UV (EtOH): λ_{max} = 231,
249, 293, 305, 333 nm.

Elementary analysis for $\text{C}_{15}\text{H}_{16}\text{N}_4$ (m.w. 252.32):

calcd. C 71.40, H 6.39, N 22.20%;

found C 71.38, H 6.63, N 22.61%.

15 Example 13

By reaction of 4-chloro-1-methylimidazo[1,2-
a]quinoxaline (example 2) with piperidine, according to a
procedure that is similar to that followed in example 3,
there is obtained 1-methyl-4-piperidinylimidazo[1,2-
20 a]quinoxaline. m.p. (DSC) = 78.9°C (onset); IR (KBr):
3018, 2927, 1502 cm^{-1} ; ^1H -NMR (CDCl_3): δ 8.05 (1H,dd), 7.65
(1H,dd), 7.5÷7.0 (3H,m), 4.25 (4H,t), 2.85 (3H,s), 1.9÷
1.6 (6H,m); UV (EtOH): λ_{max} = 249, 281, 297, 310, 329 nm.

Elementary analysis for $\text{C}_{16}\text{H}_{18}\text{N}_4$ (m.w. 266.35):

25 calcd. C 72.15, H 6.81, N 21.03%;

found C 71.84, H 7.09, N 20.70%.

Example 14

By reaction of 4-chloroimidazo[1,2-a]quinoxaline
(example 1) with morpholine, according to a procedure

that is similar to that followed in example 3, there is obtained 4-(4-morpholinyl)imidazo[1,2-a]quinoxaline. m.p. (DSC)= 142.8°C (onset); IR (KBr): 3016, 2962, 1517 cm⁻¹; ¹H-NMR (CDCl₃): δ 8.0 (1H,s), 7.8÷7.5 (3H,m), 7.5÷7.2 (2H,m), 4.4 (4H,t), 3.9 (4H,t); UV (EtOH): λ_{max} = 230, 248, 292, 304, 330 nm.

Elementary analysis for C₁₄H₁₄N₄O (m.w. 254.29):
calcd. C 66.13, H 5.55, N 22.03%;
found C 65.43, H 5.47, N 22.25%.

10 Example 15

By reaction of 4-chloroimidazo[1,2-a]quinoxaline (example 1) with N-methylpiperazine, according to a procedure that is similar to that followed in example 3, there is obtained 4-(N'-methylpiperazinyl)imidazo[1,2-a]quinoxaline, that is subsequently converted to the dihydrochloride by treatment with ethanolic HCl. m.p. (DSC)= 305.6°C (onset); IR (KBr): 2697, 1560, 1508 cm⁻¹; ¹H-NMR (DMSO-d₆/CD₃OD): δ 8.7 (1H,sb), 8.3÷8.0 (1H,m), 7.8÷7.35 (4H,m), 5.5 (4H,t), 3.7÷3.3 (4H,m), 2.8 (3H,s); UV (EtOH): λ_{max} = 230, 291, 304, 320 nm.

Elementary analysis for C₁₅H₁₇N₅·2HCl (m.w. 340.25):
calcd. C 52.95, H 5.63, N 20.58%;
found C 52.81, H 5.78, N 20.13%.

Example 16

25 By reaction of 4-chloro-1-methylimidazo[1,2-a]quinoxaline (example 2) with N-methylpiperazine, according to a procedure that is similar to that followed in example 3, there is obtained 1-methyl-4-(N'-methylpiperazinyl)imidazo[1,2-a]quinoxaline. m.p.

(DSC)= 108.0°C (onset); IR (KBr): 2928, 1535, 1510 cm^{-1} ;
 $^1\text{H-NMR}$ (CDCl_3): δ 8.1 (1H,dd), 7.7 (1H,dd), 7.4-7.0
(3H,m), 4.35 (4H,t), 2.85 (3H,s), 2.6 (4H,t), 2.3 (3H,s);
UV (EtOH): λ_{max} = 228, 248, 279, 309, 325 nm.

- 5 Elementary analysis for $\text{C}_{18}\text{H}_{19}\text{N}_3$ (m.w. 281.36):
calcd. C 68.30, H 6.81, N 24.89%;
found C 68.37, H 7.15, N 25.05%.

Example 17

- By reaction of 2,3,6-trichloroquinoxaline with
10 aminoacetaldehyde dimethyl acetal, according to a
procedure that is similar to that described in example 1,
there is obtained 8-chloroimidazo[1,2-a]quinoxalin-4(5H)-
one (m.p. >300°C). By reacting this product with 1-ethyl
propylamine according to the method described in example
15 9, there is obtained 8-chloro-4-(1-ethylpropyl-
amino)imidazo[1,2-a]quinoxaline. m.p.

- (DSC)= 125.1°C (onset); IR (KBr) : 3406, 3105, 2964, 1532
 cm^{-1} ; $^1\text{H-NMR}$ ($\text{CDCl}_3/\text{CD}_3\text{OD}$): δ 8.4 (1H,s), 8.05 (1H,d,
J=2Hz), 7.8-7.4 (3H,m), 3.85 (1H,m), 1.65 (4H,m), 0.95
20 (6H,t); UV (EtOH): λ_{max} = 230, 245, 289, 301, 326 nm.

Elementary analysis for $\text{C}_{18}\text{H}_{17}\text{ClN}_3$ (m.w. 288.78):
calcd. C 62.39, H 5.93, N 19.40%;
found C 62.17, H 6.02, N 19.46%.

Example 18

- 25 By reacting 8-chloroimidazo[1,2-a]quinoxalin-4(5H)-
one (example 17) with POCl_3 , following a procedure that
is similar to that described in example 1, there is
obtained 4,8-dichloroimidazo[1,2-a]quinoxaline that is
subsequently reacted with cyclopentylamine according to

the method described in example 3, thereby obtaining 4-cyclopentylamino-8-chloroimidazo[1,2-a]quinoxaline.

m.p.

(DSC) = 140.1°C (onset); IR (KBr): 3401, 2955, 1554 cm^{-1} ;

5 $^1\text{H-NMR}$ (CDCl_3): δ 7.8 (1H, d, $J=2\text{Hz}$), 7.7÷7.2 (4H, m), 6.15 (1H, d) 4.6 (1H, m), 2.3÷2.05 (4H, m), 2.05÷1.3 (4H, m); UV (EtOH): λ_{max} = 229, 245, 289, 301, 326, 340 nm.

Elementary analysis for $\text{C}_{15}\text{H}_{15}\text{ClN}_4$ (m.w. 286.76):

calcd. C 62.83, H 5.27, N 19.54%;

10 found C 63.04, H 5.36, N 19.64%.

Example 19

By reacting 4,8-dichloroimidazo[1,2-a]quinoxaline (example 18) with cyclohexylamine, following a procedure that is similar to that described in example 3, there is
15 obtained 4-cyclohexylamino-8-chloroimidazo[1,2-a]quinoxaline. m.p.

(DSC)=126.7°C (onset); IR (KBr): 3413, 2926, 1555 cm^{-1} ;

$^1\text{H-NMR}$ (CDCl_3): δ 7.85 (1H, s), 7.7÷7.2 (4H, m), 6.1 (1H, d),
4.2 (1H, m), 2.3÷2.0 (4H, m), 2.0÷1.2 (6H, m); UV (EtOH):
20 λ_{max} = 229, 245, 289, 301, 326, 340 nm.

Elementary analysis for $\text{C}_{16}\text{H}_{17}\text{ClN}_4$ (m.w. 300.79):

calcd. C 63.89, H 5.70, N 18.63%

found C 64.07, H 5.79, N 18.80%

Example 20

25 By reacting 2,3,6-trichloroquinoxaline with propargylamine, following a procedure that is similar to that described in example 2, there is obtained 8-chloro-1-methylimidazo[1,2-a]quinoxalin-4(5H)-one (m.p. >300°C) and, subsequently, 4,8-dichloro-1-methylimidazo[1,2-a]

quinoxaline. By reacting this product with cyclopentylamine according to the method described in example 3, there is obtained 4-cyclopentylamino-8-chloro-1-methylimidazo[1,2-a]quinoxaline. m.p.

5 (DSC) = 130.5°C (onset); IR (KBr): 3421, 2954, 1555 cm^{-1} ;
1H-NMR (CDCl_3): δ 7.95 (1H, d, J=2Hz), 7.6 (1H, d, J=9Hz),
7.35÷7.15 (2H, m), 6.1 (1H, d), 4.5 (1H, s), 2.8 (3H, s),
2.4÷2.05 (4H, m), 2.0÷1.3 (4H, m); UV (EtOH): λ_{max} = 227,
251, 276, 305, 323, 336 nm.

10 Elementary analysis for $\text{C}_{16}\text{H}_{17}\text{ClN}_4$ (m.w. 300.79):
calcd. C 63.89, H 5.70, N 18.63%
found C 63.79, H 5.79, N 18.58%

Example 21

By reacting 4,8-dichloro-1-methylimidazo[1,2-a]quinoxaline (example 20) with cyclohexylamine, following
15 a procedure that is similar to that described in example 3, there is obtained 4-cyclohexylamino-8-chloro-1-methylimidazo[1,2-a]quinoxaline. m.p.

(DSC) = 130.4°C (onset); IR (KBr): 3410, 2927, 1551 cm^{-1} ;
20 1H-NMR (CDCl_3): δ 7.95 (1H, d, J=2Hz), 7.6 (1H, d, J=9Hz),
7.4÷7.2 (2H, m), 6.1 (1H, d), 4.2 (1H, m), 2.8 (3H, 2), 2.3÷
1.1 (10H, m); UV (EtOH): λ_{max} = 227, 250, 276, 305, 323,
336 nm.

Elementary analysis for $\text{C}_{17}\text{H}_{19}\text{ClN}_4$ (m.w. 314.82):
25 calcd. C 64.86, H 6.08, N 17.80%
found C 64.86, H 6.21, N 17.91%

Example 22

By reacting 2,3,6,7-tetrachloroquinoxaline with aminoacetaldehyde dimethyl acetal, following a procedure

that is similar to that described in example 1, there is obtained 7,8-dichloroimidazo[1,2-a]quinoxalin-4(5H)-one (m.p. >300°C) and, subsequently, 4,7,8-trichloroimidazo[1,2-a]quinoxaline. By reacting this product with cyclopentylamine according to the method described in example 3, there is obtained 4-cyclopentylamino-7,8-dichloroimidazo[1,2-a]quinoxaline.

m.p. (DSC) = 139.4°C (onset); IR (KBr): 3247, 2961, 1589, 1556 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 7.8 (2H,m), 7.7 (1H,s), 7.55 (1H,s), 6.2 (1H,d), 4.6 (1H,m), 2.4÷1.3 (8H,m); UV (EtOH): λ_{max} = 235, 274, 292, 304, 330, 345 nm.

Elementary analysis for $\text{C}_{12}\text{H}_{14}\text{Cl}_2\text{N}_4$ (m.w. 321.21):
calcd. C 56.09, H 4.39, N 17.44%;
found C 56.13, H 4.41, N 17.52%.

Example 23

By reacting 4,7,8-trichloroimidazo[1,2-a]quinoxaline (example 22) with cyclohexylamine, following a procedure that is similar to that described in example 3, there is obtained 4-cyclohexylamino-7,8-dichloroimidazo[1,2-a]quinoxaline. m.p.

(DSC) = 162.3°C (onset); IR (KBr): 3332, 2929, 1587, 1550 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 7.8 (2H,m), 7.7 (1H,s), 7.55 (1H,d, $J=2\text{Hz}$), 6.2 (1H,d), 4.4 (1H,m), 2.3÷1.2 (10H,m); UV (EtOH): λ_{max} = 234, 274, 292, 304, 330, 345 nm.

Elementary analysis for $\text{C}_{16}\text{H}_{18}\text{Cl}_2\text{N}_4 \cdot 1/2\text{H}_2\text{O}$ (m.w. 344.25):
calcd. C 55.82, H 4.98, N 16.28%;
found C 55.83, H 5.00, N 16.31%

Example 24

By reacting 2,3,6,7-tetrachloroquinoxaline with

propargylamine, following a procedure that is similar to that described in example 2 there is obtained 7,8-dichloro-1-methylimidazo[1,2-a]quinoxalin-4(5H)-one (m.p. >300°C) and, subsequently, 1-methyl-4,7,8-trichloroimidazo[1,2-a]quinoxaline. By reacting this product with cyclopentylamine according to the method described in example 3, there is obtained 4-cyclopentylamino-7,8-dichloro-1-methylimidazo[1,2-a]quinoxaline.

m.p. (DSC) = 213.5°C (onset); IR (KBr): 3411, 2960, 1548 cm⁻¹; ¹H-NMR (CDCl₃): δ 8.0 (1H,s), 7.75 (1H,s), 7.25 (1H,s), 6.2 (1H,d), 4.6 (1H,m), 2.8 (3H,s), 2.3÷1.3 (8H,m); UV (EtOH): λ_{max} = 233, 279, 308, 327, 341 nm.

Elementary analysis for C₁₆H₁₄Cl₂N₄ (m.w. 335.23):

calcd. C 57.33, H 4.81, N 16.71%;

found C 57.41, H 4.82, N 16.68%.

Example 25

By reacting 2,3-dichloro-6-fluoroquinoxaline with aminoacetaldehyde dimethyl acetal, following a procedure that is similar to that described in example 1, there is obtained 8-fluoroimidazo[1,2-a]quinoxalin-4(5H)-one (m.p. >300°C) and, subsequently, 4-chloro-8-fluoroimidazo[1,2-a]quinoxaline. By reacting this product with cyclopentylamine according to the method described in example 3, there is obtained 4-cyclopentylamino-8-fluoroimidazo[1,2-a]quinoxaline.

m.p. (DSC) = 85.7°C (onset); IR (KBr) : 3255, 2964, 1551 cm⁻¹; ¹H-NMR (CDCl₃): δ 7.8 (1H,s), 7.7÷7.35 (1H,dd, J_{HF}=16Hz), 7.5 (1H,d, J=2Hz), 7.6÷7.25 (1H,dd, J_{HF}=16Hz), 7.05 (1H,dd), 6.0 (1H,d), 4.55 (1H,m), 2.35÷1.3 (8H,m);

UV (EtOH): λ_{max} = 226, 268, 285, 296, 323, 336 nm.

Elementary analysis for $\text{C}_{15}\text{H}_{15}\text{FN}_4$ (m.w. 270.31):

calcd. C 66.65, H 5.59, N 20.73%;

found C 66.36, H 5.66, N 20.86%.

5 Example 26

By reacting 4-chloro-8-fluoroimidazo[1,2-a]quinoxaline (example 25) with cyclohexylamine, following a procedure that is similar to that described in example 3, there is obtained 4-cyclohexylamino-8-fluoro-imidazo[1,2-a]quinoxaline. m.p.

(DSC) = 157.6°C (onset); IR (KBr): 3415, 2927, 1556 cm^{-1} ;
H-NMR (CDCl_3): δ 7.8 (1H, dd, $J=2\text{Hz}$), 7.7-7.35 (1H, dd, $J_{\text{HF}}=16\text{Hz}$), 7.55 (1H, d, $J=2\text{Hz}$), 7.55-7.25 (1H, dd, $J_{\text{HF}}=16\text{Hz}$), 7.05 (1H, dd), 6.0 (1H, d), 4.25 (1H, m), 2.35-1.2 (10H, m); UV (EtOH): λ_{max} = 226, 239, 285, 297, 323, 337 nm.

Elementary analysis for $\text{C}_{17}\text{H}_{17}\text{FN}_4$ (m.w. 284.33):

calcd. C 67.59, H 6.03, N 19.71%

found C 67.31, H 6.04, N 19.70%

20 Example 27

By reacting 2,3-dichloro-6,7-difluoroquinoxaline with aminoacetaldehyde dimethyl acetal, following a procedure that is similar to that described in example 1, there is obtained 7,8-difluoroimidazo[1,2-a]quinoxalin-4(5H)-one (m.p. $>300^\circ\text{C}$) and, subsequently, 4-chloro-7,8-difluoroimidazo[1,2-a]quinoxaline. By reacting this product with cyclopentylamine according to the method described in example 3, there is obtained 4-cyclopentylamino-7,8-difluoroimidazo[1,2-a]quinoxaline. m.p.

(DSC) = 146.0°C (onset); IR (KBr): 3263, 2955, 1554 cm^{-1} ;
 $^1\text{H-NMR}$ (CDCl_3): δ 7.7 (1H,s), 7.65÷7.15 (3H,m), 6.1
(1H,d), 4.6 (1H,m), 2.3÷1.3 (8H,m); UV (EtOH): λ_{max} = 225,
241, 270, 295, 323, 337 nm.

- 5 Elementary analysis for $\text{C}_{15}\text{H}_{14}\text{F}_2\text{N}_4$ (m.w. 288.30):
calcd. C 62.49, H 4.89, N 19.43%;
found C 62.46, H 5.03, N 19.65%.

Example 28

Binding on the adenosine receptors.

- 10 The binding on A_2 receptors has been carried out
according to the method described in Naunyn-Schmied.
Arch. Pharmacol. 1987, 336, 204 on preparations of
synaptosomal membranes from rat brain by incubating 200
 μg of membrane proteins for 1 h at 25°C with the
15 substance to be tested and 0.3 nM [^3H]-DPCPX in 400 μl of
50 mM Tris.HCl pH = 7.4. The non-specific binding has
been determined with 5 nM R-PIA.

- The binding on A_2 receptors has been carried out
according to the method described in FASEB J. 1989, 3,
20 A1047 on preparations of rat striatal membranes by
incubating 200 μg of membrane proteins for 1 h at 25°C
with the substance to be tested, 5 nM [^3H]-CGS21680 and
50 nM CPA. The non-specific binding has been determined
with 100 nM CPA.

- 25 The incubations were blocked by filtration by means
of a cell-harvester and, after completion of the
separation of the bound from the free, the radioactivity
contents were evaluated by liquid scintillation. The
concentration-inhibition curves were obtained by assaying

the receptor displacement at at least ten different concentrations of the test substance (all the assays were carried out in triplicate). The tested substances were dissolved in dimethyl sulphoxide and diluted in 50 mM Tris.HCl buffer pH = 7.4. The IC_{50} values have been determined by non-linear regression curves and transformed into K_i values according to the Cheng-Prusoff equation.

Table I shows the results obtained with the compounds of examples 5, 9, 13, 18, 22 and 25 of the invention.

TABLE I: Adenosine receptors affinities :

Substance	Ki, A: (nM)	Ki, A: (μM)
COMPOUND EX. 5	54	
COMPOUND EX. 9	7,9	2,5
COMPOUND EX. 13	-	2,6
COMPOUND EX. 18	23,5	
COMPOUND EX. 22	26,5	
COMPOUND EX. 25	84	

Example 29

Forced swim test

5 The test described by R.D.Porsolt et al. in Arch.Int.Pharmacodyn.Ther. 1977, 229, 327, has been carried out which is widely used as an animal model for the study of the antidepressant activity of new drugs. Male albino CD 1 mice weighing 25-35 g were used.

10 One hour before immersion in water, the test compound is intraperitoneally (i.p.) administered to the animal; the vehicle is administered to the control animals. The duration of permanence in water is 6': from the 2nd to the 6th minute the time is measured during
15 which the animal remains motionless. Table II shows the results obtained with the compounds of examples 9, 13, 16, 18 and 21 of the invention, expressed as the percent variation of the immobility time of the treated animals versus the control group. As the reference substance the
20 tricyclic antidepressant drug desipramine was used.

TABLE II: Forced swim test

Substance	Dose (mg/kg.i.p.)	%variation immobility time vs.C.
COMPOUND EX.9	0.001	-32.8***
COMPOUND EX.9	0.01	-45.3***
COMPOUND EX.13	0.1	-23.5*
COMPOUND EX.13	1	-70.7***
COMPOUND EX.16	0.1	-22.4
COMPOUND EX.16	1	-51.4***
COMPOUND EX.18	1	-42.3***
COMPOUND EX.18	10	-52.8***
COMPOUND EX.21	0.001	-33.0**
COMPOUND EX.21	0.01	-48.6***
DESIPRAMINE	7.5	-10.4**
DESIPRAMINE	30	-41.4***

(***) $p < 0.001$; (**) $p < 0.01$; (*) $p < 0.05$

Example 30

5 Tail suspension test

The test described by L.Steru, et al. in Psychopharmacology 1985, 85, 367, also widely used as an animal model for the screening of antidepressant activity has been carried out.

10 Male albino CD 1 mice weighing 25-35 g were used.

Half an hour before the test, the animal is

administered the test compound by the i.p. route; the control animals are administered the vehicle. The mouse is suspended from a horizontal bar at about 40 cm from the support plane by means of an adhesive tape applied at the end of the tail and secured to a hook. The immobility time period is registered during 6'.

Table III shows the results obtained with the compound of example 9 of the invention, as the percent change of the immobility time period of the treated animals versus the control group.

As the reference substance the tricyclic antidepressant drug desipramine was used.


TABLE III: Tail suspension test

TABLE III: Tail suspension test		
Substance	Dose (mg/kg.i.p.)	%variation immobility time vs.C.
COMPOUND EX.9	0.001	-26.7
COMPOUND EX.9	0.01	-41.1*
COMPOUND EX.9	0.1	-64.3***
DESYPRAMINE	4	-21.3
DESYPRAMINE	16	-62.4**

(***) $p < 0.001$; (**) $p < 0.01$; (*) $p < 0.05$

CLAIMS

1. A compound of the formula (I):



wherein:

10 R_1 is hydrogen or methyl;


R_2 is hydrogen, straight- or branched- chain (C_1-C_6) alkyl;

R_3 is hydrogen, straight- or branched- chain (C_1-C_6) alkyl

that is possibly substituted with OH, (C_5-C_6) cycloalkyl;

or R_2 and R_3 together form

15



wherein Z is a direct bond, or O, NR₄, R₄ being a straight- or branched- chain (C_1-C_6) alkyl;

m and n, same or different, are 1, 2 or 3;

25 R_4 and R_5 can be the same or different and are hydrogen or halogen chosen from Cl, F, Br;


and its pharmacologically acceptable salts

30 2. A compound according to claim 1 characterized in that R_1 is hydrogen or methyl, R_2 is hydrogen, R_3 is hydrogen, straight- or branched- chain (C_1-C_6) alkyl that is possibly substituted with OH, (C_5-C_6) cycloalkyl, R_2 and R_3 can be the same or different and are hydrogen,

chlorine or fluorine.


3. A compound according to claims 1 and 2 characterized in that R_1 is methyl, R_2 is hydrogen, R_3 is cyclopentyl, R_4 and R_5 are both hydrogen.

5 4. A process for the preparation of the compounds of the formula (I) according to claims 1 to 3, characterized in that a compound of formula (II)




(II)

in which R_4 and R_5 are as defined in claim 1, is reacted with amino acetaldehyde dimethyl acetal or with propargyl amine, thereby to obtain, respectively, the compounds of the formulae (III) and (IV)




(III)




(IV)

in which R_4 and R_5 are as defined in claim 1, said compounds being subsequently cyclized in an acidic milieu to give the compound of formula (V)



(V)

in which R_2 is hydrogen or methyl, R_4 and R_5 are as defined in claim 1; compound (V) is then reacted with a chlorinating agent such as $POCl_3$ to provide the compound of formula (VI)



(VI)


in which all the substituents are as defined above, which is subsequently reacted with the amine HNR_2R_3 , in which R_2 and R_3 are as defined in claim 1;

or

compound (V), prepared as described above, is reacted with a silylating agent, such as hexamethyl disilazane, and amine HNR_2R_3 , in which R_2 and R_3 are as defined in claim 1, at a temperature of between 80 and 180°C;

or

for the preparation of the compounds of formula (I) in which R_2 and R_3 are both hydrogen, the compound (VI), prepared as described above is reacted with hydrazine to give the compound of the formula (VII)



(VII)

in which R_1 , R_4 and R_5 are as defined in claim 1, which is subsequently hydrogenated according to conventional methods.

5 5. Pharmaceutical compositions characterized in that they contain at least a compound according to claim 1 as an active ingredient at an effective dosage together with one or more conventional, non-toxic excipient.

 6. Use of the compounds of claim 1 as adenosine antagonists.

10 7. Use of compounds according to claim 1 for the preparation of a medicament for the treatment of psychiatric disorders.

 8. Use of compounds according to claim 1 for the preparation of a medicament for the treatment of
15 neurological diseases of the central nervous system.

20

25

INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/IB 96/01291

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D487/04 A61K31/505 //(C07D487/04,241:00,235:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 229 452 A (WARNER ET AL.) 21 October 1980 * See Table VII, columns 31-32, Compounds no. 167-169 *	1
X	--- CHEMICAL ABSTRACTS, vol. 77, no. 13, 25 September 1972 Columbus, Ohio, US; abstract no. 88433s, SIMONOV ET AL.: "Imidazo[1,2-a]quinoxaline and its reactions" page 457; XP002024866 & KHIM. GETEROTSIKL. SOEDIN, vol. 3, 1972, pages 416-418, see abstract --- -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

10 February 1997

Date of mailing of the international search report

10.03.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 96/01291

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 780 464 A (TRIVEDI) 25 October 1988 see column 1 - column 5-8 ---	1
Y	JOURNAL OF MEDICINAL CHEMISTRY, vol. 34, no. 3, 1991, pages 1202-1206, XP002024865 P.J.M. VAN GALEN ET AL.: "1H-Imidazo[4,5-c]quinolin-4-amines: Novel Non-Xanthine Adenosine Antagonists" cited in the application see the whole document ---	1
A	US 5 182 386 A (ALBAUGH ET AL.) 26 January 1993 cited in the application see column 1-6 ---	1
A	WO 94 22865 A (BASF) 13 October 1994 cited in the application see page 1-2 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 96/01291

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4229452	21-10-80	AU-B- 521807	29-04-82
		AU-A- 3233178	19-07-79
		BE-A- 862608	03-07-78
		CA-A- 1149385	05-07-83
		CA-A- 1164877	03-04-84
		CH-A- 636102	13-05-83
		CH-A- 640229	30-12-83
		DE-A- 2800480	13-07-78
		DE-C- 2858766	05-04-90
		FR-A- 2376858	04-08-78
		GB-A- 1579498	19-11-80
		GB-A- 1579499	19-11-80
		JP-B- 1030825	22-06-89
		JP-C- 1545999	28-02-90
		JP-A- 61171470	02-08-86
		JP-C- 1450207	11-07-88
		JP-A- 53087398	01-08-78
		JP-B- 62059110	09-12-87
		LU-A- 78813	18-09-78
		NL-A- 7800114	11-07-78
		SE-B- 436359	03-12-84
		SE-A- 7800138	08-07-78
		SE-B- 460477	16-10-89
		SE-A- 8300945	21-02-83
		US-A- 4225724	30-09-80
		US-A- 4236015	25-11-80
		US-A- 4198508	15-04-80
		US-A- 4191766	04-03-80
		US-A- 4191767	04-03-80
		US-A- 4172947	30-10-79
		US-A- 4160097	03-07-79
		US-A- 4200750	29-04-80
		US-A- 4197403	08-04-80

US-A-4780464	25-10-88	NONE	

US-A-5182386	26-01-93	AU-A- 2546792	16-03-93
		WO-A- 9304066	04-03-93
		US-A- 5473073	05-12-95

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter. Application No

PCT/IB 96/01291

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9422865	13-10-94	DE-A- 4329970	06-10-94
		AU-A- 6428294	24-10-94
		BR-A- 9406034	02-01-96
		CA-A- 2158167	13-10-94
		CN-A- 1120336	10-04-96
		CZ-A- 9502464	13-03-96
		EP-A- 0691970	17-01-96
		FI-A- 954614	28-09-95
		HR-A- 940210	31-12-96
		JP-T- 8508271	03-09-96
		NO-A- 953892	29-09-95
		PL-A- 310950	08-01-96
		SI-A- 9420023	30-04-96
		ZA-A- 9402246	02-10-95
